

Composition:

Each kg powder contains tylosin tartrate 625g



Indications:

Poultry: Prevention and Treatment of CRD, infectious Coryza, Staphylococcosis and as immunity enhancer.

Prevention and control of mycoplasma infections

Dosage : Prophylactic use:

Per 1000 birds: Orally through drinking water in two divided doses.

Broilers : First 3 days: 8 grams/day

22nd or 23rd or 24th day: 72 grams / day (only one dose).

Layers : First 3 days: 8 grams/day 25th day: 40 grams / day 70th day: 120grams / day 125th day: 192 grams / day

Broiler Breeders :

4th to 7th day: 11.2 grams/ day (4 days)

13th to 14th day: 11.4 grams/ day (2 days)

20th to 21st day: 56 grams/ day (2 days)

26th to 27th day: 72 grams/ day (2 days)

34th to 35th day: 88 grams/ day (2 days)

66th day: 152 grams/ day (1 day)

87th day: 200 grams/ day (1 day)

140th day: 264 grams/ day (1 day)

Layer Breeders :

4th to 7th day : 9.6 grams/ day (4 days)

13th to 14th day : 20 grams/ day (2 days)

20th to 21st day : 40 grams/ day (2 days)

26th to 27th day: 53 grams/ day (2 days)

34th to 35th day : 66 grams/ day (2 days)

64th day : 80 grams/ day (1 day)

85th day : 160 grams/ day (1 day)

140th day : 232 grams/ day (1 day)

(or) as advised by a veterinarian.

Special note: The above doses are calculated at the rate of 176 mg/Kg. body weight. The dosage may be increased or decreased as per the body weights.

Packing : 50 gm, 100 gm, 200 gm, 500 gm, 1 kg. 5kg

Manufacturer: Hebei Weierli Animal Pharmaceutical Group Co., Ltd

Exporter: Hebei Nuob Trade Co., Ltd

Website: www.victorypharm.com

Alibaba: www.victorypharm.en.alibaba.com

Email: julia@victorypharm.com

Mobie&Whatsapp: +86 13191871625

Tel: +86-311-89265710 **Fax:** +86-311-58052865