

Description

Toltrazuril is an anticoccidial with activity against Eimeria spp. in poultry:

- Eimeria acervulina, brunetti, maxima, mitis, necatrix and tenella in chickens.
- Eimeria adenoides, gallopilonis and meleagrimitis in turkeys.



Composition

Each ml Contains: Toltrazuril 25 mg.

Indications

Coccidiosis of all stages like schizogony and gametogony stages of Eimeria spp. in chickens and turkeys.

Contra indications

Administration to animals with impaired hepatic and/or renal function.

Side effects

At high dosages in laying hens egg-drop and in broilers growth inhibition and polyneuritis can occur.

Dosage

For oral administration:

- 500 ml per 500 litre of drinking water (25 ppm) for continuous medication over 48 hours, or
- 1500 ml per 500 litre of drinking water (75 ppm) given for 8 hours per day, on 2 consecutive days

This corresponds to a dose rate of 7 mg of toltrazuril per kg of body weight per day for 2 consecutive days.

Note: supply the medicated drinking water as the only source of drinking water. Do not administer to poultry producing eggs for human consumption.


Withdrawal times

For meat:

- Chickens : 18 days.
- Turkeys : 21 days.

Packaging

Bottle of 100, 500 and 1000 ml.





Manufacturer: Hebei Pude Animal Medicine Co., Ltd

Exporter: Hebei Nuob Trade Co., Ltd

Website: www.victorypharm.com

Alibaba: www.victorypharm.en.alibaba.com

Email: julia@victorypharm.com

Mobie&Whatsapp: +86 13191871625

Tel: +86-311-89265710 **Fax:** +86-311-58052865



Address: B-1601, Lechen Business Building, No.260 Huaian Road, Shijiazhuang City, Hebei, China