

# **维尔利集团** **Amoxa 100 WSP** **WEIERLI GROUP** Amoxicillin Water Soluble Powder 10%

Amoxicillin is a semi-synthetic penicillin possessing a broad spectrum of antimicrobial activity. It acts bactericidal against a number of Gram positive and negative microorganism, particularly against E. coli, Streptococcus spp., Pasteurella spp. salmonella spp. Bordetella bronchiceptica, Staphylococcus and others.

## Composition

Each Kg contains Amoxicillin Trihydrate 100 g (Act)



## Indication

Treatment of the disease caused by the following micro-organism susceptible to amoxicillin; Staphylococcus spp., Streptococcus spp., Pasteurella spp., Escherichia coli, Hemophilus spp., Actinobacillus pleuropneumoniae.

Calf (less than 5 months old): pneumonia, diarrhea caused by Escherichia coli

Swine: pneumonia, diarrhea caused by Escherichia coli

## Dosage & Administration

The following dosage is mixed with feed or drinking water and orally administer once or twice a day. (However, do not take more than 5 days)

	Indication	Daily Dosage of this drug/1kg of b.w.	Daily Dosage of Amoxicillin /1kg of b.w.
Calves	Pneumonia	30-100 mg	3-10 mg
	Diarrhea caused by	50-100 mg	5-10 mg
	Escherichia coli		
Swine	Pneumonia	30-100 mg	3-10 mg



Poultry: general dosage is 10mg amoxicillin per kg b.w per day.

Prevebtion: 1g per 2 liter of drinking water, continue for 3 to 5 days

Treatment: 1g per 1 liter of drinking water, continue for 3 to 5 days

## Packaging

500g, 1kg, 5kg, 10kg, 25k



## Storage and expiry date

Store in an airtight container at dry room temperature (1 to 30o C) protected from light.  
24 months from date of manufacture

## Precaution

- A. Do not use for animals with shock and hypersensitive response to this drug.
- B. Side effect
  - 1. Penicillin antibiotics may cause diarrhea by inhibiting the intestinal normal bacterial flora and bring about abdominal pain by gastroenteritis or colitis, digestive system abnormalities such as anorexia, watery diarrhea or hemafecia, nausea and vomiting and etc.
  - 2. Penicillin antibiotics may induce nervous system abnormalities such as convulsions and seizures and hepatotoxicity when overdosing
- C. Interaction
  - 1. Do not administer with macrolide (erythromycin), aminoglycoside, chloramphenicol, and tetracycline antibiotics.
  - 2. Gentamicin, bromelain and probenecid may increase the efficacy of this drug.
- D. Administration for pregnant, lactating, newborn, weanling and debilitated animals :  
Do not administer to laying chickens
- E. Usage note
  - 1. When administering by mixing with feed or drinking water, mix homogeneously to prevent from drug accident and to achieve its efficacy.
  - 2. Withdrawal period
    - Poultry: 7 days
    - Cattle: 10 days
    - Swine: 10 days

## F. Precaution on storage

- ① To prevent safety accidents, keep out of reach of children and an animal.
- ② Observe the preservation instruction since it may bring about a change in effectiveness and stability.
- ③ Dispose the expired products without using it.
- ④ Use as soon as possible after opening, the rest of it should be sealed in the original packaging container and store in a dry place protected from light.
- ⑤ Do not use used containers or wrapping paper for other purposes and discard it safely.
- ⑥ Keep out of reach of the children.

## G. Other Precaution

- 1, It is for animal use, so never use it for human.
- 2, Consult with your veterinarian.
- 3, Use after reading the precaution note.





- 4, Since safety and effectiveness other than the designated animal has not been established, do not use it arbitrarily
- 5, Record all usage history for prevention of abuse and tolerance appearance
- 6, Abuse and misuse may bring about economic loss such as drug accidents and remaining animal food residues, observe the dosage & administration.
- 7, Abuse and misuse may bring about economic loss such as drug accident, so do not administer with drugs containing the same ingredients concurrently.
- 8, Do not administer it with the drug contains same ingredients concurrently.
- 9, If you are not in compliance with withdrawal period, that may bring about remaining drugs in animal foods, so accurately calculate and comply with withdrawal period after calculation time
- 10, Wear gloves, masks, protective equipment during handling to avoid skin contact and inhalation.
- 11, Consult with doctor as soon as abnormality is found.
- 12, When administering by mixing with feed or drinking water, mix homogeneously to prevent from drug accident and to achieve its efficacy.



**Manufacturer:** Hebei Weierli Animal Pharmaceutical Group Co., Ltd

**Exporter:** Hebei Nuob Trade Co., Ltd

**Website:** [www.victorypharm.com](http://www.victorypharm.com)

**Alibaba:** [www.victorypharm.en.alibaba.com](http://www.victorypharm.en.alibaba.com)

**Email:** [julia@victorypharm.com](mailto:julia@victorypharm.com)

**Mobie&Whatsapp:** +86 13191871625

**Tel:** +86-311-89265710 **Fax:** +86-311-58052865

**Address:** B-1601, Lechen Business Building, No.260 Huaian Road, Shijiazhuang City, Hebei, China